

ANKK1/DRD2

Clinical Utility

Dopamine, a key neurotransmitter that controls cognition, emotion, locomotor activity, and other endocrine functions, exerts its action by binding to five different receptors, including the dopamine D2 receptor (DRD2). Dysregulation of dopaminergic signal transmission is found in many pathological conditions such as Parkinson's disease and schizophrenia, and compounds that act as DRD2 agonists or antagonists are used to treat these conditions. Therapeutic and adverse events of several antipsychotics both result from their high affinity to antagonize DRD2.

Assay Interpretation

Within the several genetic variants of DRD2 that are relevant to disease susceptibility and therapeutic response, the Taq1A (32806C>T; rs1800497) is one of the most studied. This variant is located downstream of the DRD2 gene within the ankyrin repeat of the ANKK1 gene. The presence of the Taq1A T variant defines the A 1 allele that is associated with a reduced DRD2 gene expression and function. The A2 allele defines the reference allele. The frequency of the minor Taq1A T allele differs among ethnic populations. It occurs in 22% of Caucasians, and 42% of Asians and Africans.

The reference range for the Taq1A variant is 32806C>T CC (A2/A2) and is associated with a normal DRD2 expression.

Clinical Implications

The presence of the Taq 1A A 1 allele (32806C>T) seems to be associated with nicotine dependence and the efficacy of bupropion and nicotine replacement therapy. Smokers carrying the normal DRD2 phenotype (A2/A2 genotype) using bupropion for smoking cessation are three times more likely to be abstinent at the end of treatment than non-carriers of this genotype. Smokers with the Taq1A T variant allele (A1) seem to derive greater benefits from nicotine replacement therapies. Antipsychotic agents have been associated with hyperprolactinemia and tardive dyskinesia (TD). TD-positive patients taking antipsychotics have a higher A2 allele frequency, while A1 allele is overrepresented among those experiencing hyperprolactinemia.

References

1: Young et al. Prolactin levels in antipsychotic treatment of patients with schizophrenia carrying the DRD2*A1 allele. Br J Psychiatry. 2004 Aug;185:147-51. 2: Zai et al. Meta-analysis of two dopamine D2 receptor gene polymorphisms with tardive dyskinesia in schizophrenia patients. Mol Psychiatry. 2007 Sep;12(9):794-5. 3: Bakker et al. Antipsychotic-induced tardive dyskinesia and polymorphic variations in COMT, DRD2, CYP1A2 and MnSOD genes: a meta-analysis of pharmacogenetic interactions. Mol Psychiatry. 2008 May;13(5):544-56. 4: Johnstone et al. Pharmacogenetics. 2004 Feb;14(2):83-90. Genetic variation in dopaminergic pathways and short-term effectiveness of the nicotine patch. 5: David et al. Bupropion efficacy for smoking cessation is influenced by the DRD2 Taq1A polymorphism: analysis of pooled data from two clinical trials. Nicotine Tob Res. 2007 Dec;9(12):1251-7.

APOLIPOPROTEIN E

Clinical Utility

Apolipoproteins (APO) are structural constituents of lipoprotein particles that have critical roles in blood lipid metabolism and transport. Apolipoprotein E (APOE) is a major constituent of triglyceride-rich chylomicrons, very low-density lipoproteins (VLDL), and some subclasses of high-density lipoproteins (HDL). The primary function of APOE is to transport cholesterol from the cells in the blood vessel wall to the liver for excretion. Defects in apolipoprotein E (APOE) can result in dyslipidemia, which is an important risk factor in the genesis of atherosclerosis and subsequent development of cardiovascular disease (CVD).

Assay Interpretation

There are three common APOE alleles designated E2, E3, and E4, resulting from combinations of the two genetic polymorphisms 388T>C and 526 C>T. These alleles result in E2, E3, and E4 protein isoforms, respectively. The approximate allele frequencies for most populations are 8- 12% for E2, 74-78% for E3, and 14-15% for E4.

The reference ranges for both mutations of APOE are 388TT and 526CC. This is consistent with a E3/E3 genotype and a normal APOE function.

Clinical Implications

- The APOE E3/E3 genotype is considered the normal genotype and is associated with normal lipid metabolism. It is not associated with an increased risk of atherosclerotic CVD.
- The APOE E2 allele is strongly associated with type III hyperlipoproteinemia. This may increase the risk for premature CVD.
- Patients with symptoms (xanthomas) and with a lipid profile consistent with type III hyperlipidemia are candidates for APOE genotype analysis. Over 90% of individuals presenting the type III hyperlipoproteinemia have the rare E2/E2 genotype. However, only 1-5% of individuals with this genotype develop type III hyperlipoproteinemia, suggesting that other genetic, hormonal, or environmental factors must contribute to this disease.
- Although individuals with the APOE E2/E2 genotype are at higher risk of premature vascular disease, they may never develop the disease because this genotype is only one of the risk factors.
- In normolipidemic patients, the E2 allele is associated with lower serum cholesterol concentrations, and may confer a protection against hypercholesterolemia.
- The APOE E2/E4 genotype is associated with type III hyperlipoproteinemia in patients who are also heterozygous for familial hypercholesterolemia.
- The APOE E4 allele has been linked to pure elevations of low-density lipoproteins (LDL), and the E4/E4 and E3/E4 genotypes are associated with increased serum cholesterol levels and increased risk of CVD.

References

1: Eichner et al. Apolipoprotein E polymorphism and cardiovascular disease: a HuGE review. *Am J Epidemiol.* 2002 Mar 15;155(6):487-95. 2 : Koch et al. Apolipoprotein E gene epsilon2/epsilon3/epsilon4 polymorphism and myocardial infarction: case-control study in a large population sample. *Int J Cardiol.* 2008 Mar 28;125(1):116-7. 3: Hanis et al. Effects of the apolipoprotein E polymorphism on levels of lipids, lipoproteins, and apolipoproteins among Mexican-Americans in Starr County, Texas. *Arterioscler Thromb.* 1991 Mar-Apr;11(2):362-70. 4 : Klos et al. Linkage analysis of plasma ApoE in three ethnic groups: multiple genes with context-dependent effects. *Ann Hum Genet.* 2005 Mar;69(Pt 2):157-67. 5 : Bennet et al. Association of apolipoprotein E genotypes with lipid levels and coronary risk. *JAMA.* 2007 Sep 19;298(11):1300-11. 6 : Ciftcioglu et al. The association of apolipoprotein E polymorphism and lipid levels in children with a family history of premature coronary artery disease. *J Clin Lipidol.* 2012 Jan-Feb;6(1):81-7. 7 : Kofler et al. Apolipoprotein E genotype and the cardiovascular disease risk phenotype: impact of sex and adiposity (the FINGEN study). *Atherosclerosis.* 2012 Apr;221(2):467-70. 8 : Carvalho-Wells et al. Interactions between age and apoE genotype on fasting and postprandial triglycerides levels. *Atherosclerosis.* 2010 Oct;212(2):481-7. 9 : Sima et al. Apolipoprotein E polymorphism-a risk factor for metabolic syndrome. *Clin Chem Lab Med.* 2007;45(9):1149-53. 10 : Graner et al. Apolipoprotein E polymorphism is associated with both carotid and coronary atherosclerosis in patients with coronary artery disease. *Nutr Metab Cardiovasc Dis.* 2008 May;18(4):271-7.

COMT

Clinical Utility

Catechol-O-Methyltransferase (COMT) is an enzyme responsible for the metabolism of catecholamines and catechol-estrogens in the central nervous system and other organs. Dopamine is cleared mainly by COMT in the frontal cortex, and a reduced activity of this enzyme results in higher synaptic levels of dopamine, which affects prefrontal cortex cognitive response to certain drugs. A single nucleotide polymorphism of the COMT gene produces an amino acid change from valine to methionine (Val158Met) and reduces the enzyme activity by 3- to 4-fold.

Assay Interpretation

The most well studied COMT genetic variant (rs4680; 472G>A) is the one resulting in a valine to methionine change at codon 158. The variant allele called the Met allele is found in 30-47% of Caucasians, 23% of Africans, and 27-32% of Asians. The phenotype is defined by the presence of the reduced activity Met allele (A variant). The wild-type genotype (Val/Val; GG) predicts a high/normal COMT activity, the heterozygous genotype (Val/Met; GA) predicts an intermediate COMT activity, and the homozygous (Met/Met; M) results in a low COMT activity.

The reference range for COMT metabolic status is COMT Val158Met GG (Val/Val) (wild-type), which is consistent with a high/normal COMT activity.

Clinical Implications

Several complex associations with the Val158Met variant as a risk factor for numerous diseases have been found, but seem to have limited predictive value. The response to some psychotropic medications seems to be dependent to some extent upon the COMT status. In general, the wild-type genotype result predicts a good response to methylphenidate and amphetamines in the treatment of attention deficit hyperactivity disorder. In the treatment of pain, patients who are homozygous for the Met allele require lower doses of morphine to achieve analgesia.

References

1: De Gregori et al. Genetic variability at COMT but not at OPRM1 and UGT2B7 loci modulates morphine analgesic response in acute postoperative pain. *Eur J Clin Pharmacol.* 2013 May 19. 2 : Hamidovic et al. Catechol-O-methyltransferase val158met genotype modulates sustained attention in both the drug-free state and in response to amphetamine. *Psychiatr Genet.* 2010 Jun;20(3):85 -92. 3 : Blasi et al. Effect of catechol-O-methyltransferase val158met genotype on attentional control. *J Neurosci.* 2005 May 18;25(20):5038-45. 4 : Mattay et al. Catechol O-methyltransferase val158-met genotype and individual variation in the brain response to amphetamine. *Proc Natl Acad Sci U S A.* 2003 May 13;100(10):6186-91.

CYP1A2

Clinical Utility

The cytochrome P450 1A2 (CYP1A2) accounts for 13% of total CYP in the human liver, and is responsible for metabolizing 8-10% of commonly used drugs as well as natural compounds such as caffeine. A large inter-individual variability in the elimination of drugs that are metabolized by CYP1A2 has been observed, which has been ascribed to genetic variations and environmental factors. CYP1A2 activity is highly inducible (increased) by environmental factors including smoking (tobacco), some drugs, and several dietary compounds (cruciferous vegetables) .

Assay Interpretation

More than 20 different alleles have been characterized for the CYP1A2 gene, and some have been shown to affect enzyme activity and its sensitivity towards inducers (inducibility). The CYP1A2*1F is the most studied allele and results in a rapid metabolizer phenotype in the presence of inducers, while CYP1A2*1 K and *1C alleles result in enzymes that are less active and less sensitive to induction. The CYP1A2*1F allele is found in 25-50% of Caucasians, 30% of Asians, and 50% of Ethiopians. The genotype-phenotype relationship for CYP1A2 is not well established, and can be expressed in terms of metabolic capacity as well as sensitivity towards induction (inducibility). Individuals are predicted to have CYP1A2 normal, intermediate, or poor metabolic capacity, with high, possible, or low inducibility depending on their genotype.

The reference range for CYP1A2 metabolic status is CYP1A2 *1A/ *1A, which is consistent with a normal metabolizer that is possibly inducible.

Clinical Implications

CYP1A2 genotype can help identify patients with high or low sensitivity to inducing agents, especially those released during smoking. **The clinical relevance of this sensitivity becomes important in patients who are smokers or who quit smoking.** Patients with the highly inducible genotype CYP1A2*1F/*1F can experience loss of response to drug substrates while they are exposed to dietary or environmental inducers, and therefore may require higher doses.

The following drugs used in the management of pain and various psychiatric conditions are metabolized extensively by CYP1A2 and are sensitive to its function: clozapine (Clozaril), duloxetine (Cymbalta), olanzapine (Zyprexa), and tizanidine (Zanaflex). CYP1A2 also metabolizes other important drugs such as melatonin, ondansetron (Zofran), pomalidomide (Pomalyst), ramelteon (Rozerem), ropivacaine (Naropin), and tasimelteon (Hetlioz).

CYP1A2 metabolism is highly sensitive to inhibition and induction, and the occurrence of drug-drug interactions can have profound effects on the pharmacokinetics, response, and safety profiles of many CYP1A2 drug substrates.

Inhibitors

Some known **strong** CYP1A2 inhibitors include: ciprofloxacin (Cipro), enoxacin (Penetrex), fluvoxamine (Luvox), and zafirlukast (Accolate). Some known **moderate to weak** CYP1A2 inhibitors include: oral contraceptives, mexiletine (Mexitil), allopurinol (Zyloprim), peginterferon alfa- 2a (Pegasys), norfloxacin (Norflox), ticlopidine (Ticlid), vemurafenib (Zelboraf), and zileuton (Zyflo).

Inducers

Known CYP1A2 inducers include: carbamazepine (Tegretol), montelukast (Singulair), rifampin (Rifadin), phenytoin (Dilantin), moricizine (Ethmozine), omeprazole (Prilosec), phenobarbital, and primidone (Mysoline). Some dietary and environmental compounds found in cigarette smoke, cruciferous vegetables, and charcoal-grilled food can also increase CYP1A2 activity.

References

1: Metabolic Drug Interactions. RH Levy, KE Thummel, WF Trager. Publisher: Lippincott Williams & Wilkins (March 15, 2000). 2: Zhou et al. Polymorphism of human cytochrome P450 enzymes and its clinical impact. Drug Metab Rev. 2009;41(2):89-295 3 : Thom et al. PharmGKB summary: very important pharmacogene information for CYP1A2. Pharmacogenet Genomics. 2012 Jan;22(1):73-7. 4 : Akiillu et al. Genetic polymorphism of CYP1A2 in Ethiopians affecting induction and expression: characterization of novel haplotypes with single-nucleotide polymorphisms in intron 1. Mol Pharmacol. 2003 Sep;64(3):659-69. 5 : Zhou et al. Structure, function, regulation and polymorphism and the clinical significance of human cytochrome P450 1A2. Drug Metab Rev. 2010 May;42(2):268-354.

Genemarkers Gene Monographs

126 E. South Street | Kalamazoo | Michigan | 49007

CYP2B6

Clinical Utility

The cytochrome P450 2B6 (CYP2B6) is involved in the metabolism of 4% of clinically important medications. This enzyme is highly polymorphic: to date, 37 different variants have been identified. The CYP2B6 assay identifies some common variants that are associated with variability in enzyme activity.

Assay Interpretation

CYP2B6 enzyme activity defines a normal or an abnormal (intermediate or poor) metabolizer status for a given individual. Several variant alleles have been identified and result in different CYP2B6 isoforms that functionally are fully active, partially active, inactive, or have increased activity. The CYP2B6*1 allele is considered wild-type and encodes a functionally active enzyme (normal). The alleles *6, *7, *9, *11, *16, *18, and *36 encode a decreased activity enzyme. The *22 allele represents a gain-of-function allele. The functional impact of variants that define the CYP2B6 *4 and *5 alleles is drug dependent.

The most common functionally deficient allele is CYP2B6*6. It is found in 7-18%, 10-17%, 23%, and 33% of Caucasians, Asians, Mexican-Americans, and African-Americans, respectively. CYP2B6 *18 is found only in individuals of African descent, with a frequency of 4-7%.

The genotype-phenotype relationship is not well established, and there is a lack of consistency regarding the clinical impact of certain allelic variants. The following provisional genotype-to-phenotype assignment can be used: individuals with two fully active alleles are considered normal (extensive) metabolizers. Individuals with one fully active allele with either a partially active or an inactive allele are considered intermediate metabolizers. Individuals with two partially active alleles or two inactive alleles are considered poor metabolizers. Individuals carrying two increased function alleles or one active allele with a gain-of-function allele are classified as normal/rapid metabolizers.

The reference range for CYP2B6 metabolic status is CYP2B6 *1/*1, which is consistent with a normal metabolizer.

Clinical Implications

CYP2B6 plays a role in the metabolism of the following drugs: artemisinin, bupropion (Wellbutrin), cyclophosphamide (Cytoxan), efavirenz (Sustiva), ketamine (Ketalar), meperidine (Demerol), methadone (Dolophine), nevirapine (Viramune), propofol (Diprivan), and selegiline (Eldepryl).

The impact of CYP2B6 polymorphism on the pharmacokinetics and the clinical response have been studied in patients taking methadone, bupropion, and efavirenz. Limited evidence exists regarding the clinical impact of other polymorphisms.

Inhibitors or inducers of the CYP2B6 enzyme may modify its activity and change the patient's metabolizer status. This can result in drug-drug interactions when a drug substrate is prescribed with known CYP2B6 inhibitors or inducers.

Inhibitors

Some known CYP2B6 inhibitors include: clopidogrel, darunavir, prasugrel, ticlopidine, voriconazole, ritonavir, and thiotepa.

Inducers

Some CYP2B6 inducers include: artemether, carbamazepine, dabrafenib, efavirenz, metamizole, nevirapine, phenobarbital, phenytoin, rifampin, ritonavir, and St. John's wort.

References

1: CYP2B6 Allele Nomenclature: www.cypallele.ki.se/cyp2b6.htm 2: Li et al. Worldwide variation in human drug-metabolism enzyme genes CYP2B6 and UGT2B7: implications for HIV/AIDS treatment. *Pharmacogenomics*. 2012 Apr ;13(5):555-70. 3: Li et al. The CYP2B6*6 Allele Significantly Alters the N-Demethylation of Ketamine Enantiomers In Vitro. *Drug Metab Dispos*. 2013 Jun;41(6):1264-72. 4: Zanger and Klein. Pharmacogenetics of cytochrome P450 2B6 (CYP2B6): advances on polymorphisms, mechanisms, and clinical relevance. *Front Genet*. 2013;4:24. 5: Zanger et al. Polymorphic CYP2B6: molecular mechanisms and emerging clinical significance. *Pharmacogenomics*. 2007 Jul;8 (7):743-59. 6: Zhu et al. CYP2B6 and bupropion's smoking-cessation pharmacology: the role of hydroxybupropion. *Clin Pharmacol Ther*. 2012 Dec;92(6):771-7. 7: Benowitz et al. Influence of CYP2B6 genetic variants on plasma and urine concentrations of bupropion and metabolites at steady state. *Pharmacogenet Genomics*. 2013 Mar;23(3):135-41.

Genemarkers Gene Monographs

126 E. South Street | Kalamazoo | Michigan | 49007

CYP2C9

Clinical Utility

The cytochrome P450 2C9 (CYP2C9) is involved in the metabolism of 15% of clinically important medications. This enzyme is highly polymorphic: to date, 30 variants have been identified. The CYP2C9 assay identifies some common variants that are associated with variability in CYP2C9 enzyme activity, which has important pharmacological and toxicological implications for anticonvulsants, anticoagulants, and certain antidiabetics.

Assay Interpretation

CYP2C9 enzyme activity defines a normal or abnormal (intermediate and poor) metabolizer status for a given individual. Several variant alleles have been identified and result in different CYP2C9 isoforms that functionally are fully active, partially active, or inactive. The CYP2C9*1 allele is considered wild-type and encodes a functionally active enzyme (normal). The alleles *2, *3, *4, *5, and *11 encode a partially active enzyme. The allele *6 is a null (inactive) allele corresponding to a whole gene deletion.

The genotype-phenotype relationship is established based on the allele's activity. Individuals with two fully active alleles are considered normal (extensive) metabolizers. Individuals with one fully active allele with either a partially active or an inactive allele are considered intermediate metabolizers. Individuals with two partially active alleles or with two inactive alleles are considered poor metabolizers.

The reference range for CYP2C9 metabolic status is CYP2C9 *1r1, which is consistent with a normal metabolizer.

Clinical Implications

Abnormal CYP2C9 activity affects the therapeutic outcome of a variety of drugs used to treat cardiovascular and other conditions. Following the administration of drug substrates, the clinical manifestation in a poor or an intermediate metabolizer depends on the characteristics of the drug (i.e., the amount of drug/metabolites that is cleared by the enzyme), and the safety and pharmacological profiles of the drug and its metabolites. Within the medications used to treat cardiovascular conditions, there is compelling evidence that the response to certain angiotensin II inhibitors, statins, and anticoagulants is altered in individuals exhibiting abnormal CYP2C9 activity.

CYP2C9 plays a role in the metabolism of the following psychotropic drugs: fluoxetine (Prozac), phenytoin (Dilantin), and primidone (Mysoline). Several NSAIDs and Cox-2 inhibitors are substrates of CYP2C9, and patients with reduced CYP2C9 activity may have higher plasma levels of celecoxib (Celebrex), flurbiprofen (Ocufen), piroxicam (Feldene), or meloxicam (Mabie). CYP2C9 plays a minor role in the elimination of diclofenac (Voltaren), sulindac (Clinoril), and naproxen (Aleve).

Cardiovascular medications that are metabolized by CYP2C9 include warfarin (Coumadin), fluvastatin (Lescol), losartan (Cozaar), and irbesartan (Avapro).

Other important drugs metabolized by CYP2C9 include antidiabetics such as tolbutamide, glibenclamide (Micronase), glipizide (Glucotrol), and nateglinide (Starlix).

Inhibitors or inducers of the CYP2C9 enzyme may modify its activity and change the patient's metabolizer status. This can result in drug-drug interactions when a drug substrate is prescribed with known CYP2C9 inhibitors or inducers.

Inhibitors

Some known CYP2C9 inhibitors include: amiodarone (Cordarone), 5-fluorouracil (Adrucil), chloramphenicol, cimetidine (Tagamet), danazol (Danocrine), disulfiram (Antabuse), fluconazole (Diflucan), fluoxetine (Prozac), fluvoxamine (Luvox), miconazole (Oravig), oxandrolone (Oxandrin), capecitabine (Xeloda), co-trimoxazole (Septra), delavirdine (Rescriptor), etravirine (Intelence), fluvastatin (Lescol), efavirenz (Sustiva), gemfibrozil (Lipid), lomitapide (Juxtapid), metronidazole (Flagyl), phenytoin (Dilantin), sulfamethoxazole (Bactrim), sulfinpyrazone (Anturane), tamoxifen (Nolvadex), toremifene (Fareston), tigecycline (Tygacil), voriconazole (Vfend), and zafirlukast (Accolate).

Inducers

Some known CYP2C9 inducers include: carbamazepine (Tegretol), rifampin (Rifadin, Rimactane), rifapentine (Priftin), St. John's wort, enzalutamide (Xtandi), aprepitant (Emend), bosentan (Tracleer), dabrafenib (Tafinlar), phenobarbital, primidone (Mysoline), phenytoin (Dilantin), and ritonavir (Norvir).

References

Swen et al. Pharmacogenetics: from bench to byte-an update of guidelines. Clin Pharmacol Ther. 2011 May;89(5):662-73. 2: Wilffert et al. KNMP working group Pharmacogenetics. From evidence based medicine to mechanism based medicine. Reviewing the role of pharmacogenetics. Int J Clin Pharm. 2011 Feb;33(1):3-9. 3: Wang et al. Genetic polymorphism of the human cytochrome P450 2C9 gene and its clinical significance. Curr Drug Metab. 2009 Sep;10(7):781-834. 4- Wyatt et al. Pharmacogenetics of nonsteroidal anti-inflammatory drugs. Pharmacogenomics J. 2012 Dec;12(6):462-7

CYP2C19

Clinical Utility

The cytochrome P450 2C19 (CYP2C19) is involved in the metabolism of 10% of clinically important medications. This enzyme is highly polymorphic: more than 30 different variant alleles have been identified. The CYP2C19 assay identifies some common variants that are associated with variability in CYP2C19 enzyme activity, which has important pharmacological and toxicological implications for antidepressants, benzodiazepines, antiplatelets, and proton-pump inhibitors.

Assay Interpretation

CYP2C19 enzyme activity defines a normal or abnormal (intermediate, poor, or rapid) metabolizer status for a given individual. Several variant alleles have been identified and result in different isoforms of the CYP2C19 enzyme that functionally are fully active, partially active, inactive, or with increased activity. The CYP2C19*1 allele is considered wild-type and encodes a functionally active enzyme (normal). The alleles *2, *3, *4, *5, *6, and *8 encode an inactive enzyme and are referred to as loss-of-function alleles. Individuals with a *17 allele have an increased CYP2C19 activity.

The genotype-phenotype relationship is established based on the allele's activity. Individuals with two fully functional alleles are considered normal (extensive) metabolizers. Individuals with one or two loss-of-function alleles are considered intermediate or poor metabolizers, respectively. Individuals with one or two increased function alleles are considered rapid or ultra-rapid metabolizers, respectively. Because of limited evidence, an individual with one increased function allele and one loss-of-function allele is provisionally classified as an intermediate metabolizer.

The reference range for CYP2C19 metabolic status is CYP2C19 *1/ *1, which is consistent with a normal metabolizer.

Clinical Implications

There is substantial evidence linking the CYP2C19 polymorphisms to variability in the pharmacological and safety profiles of the following therapies used in psychiatric conditions and pain management: amitriptyline (Elavil), sertraline (Zoloft), clobazam (Onfi), citalopram (Celexa), escitalopram (Lexapro), diazepam (Valium), imipramine (Tofranil), and carisoprodol (Soma). CYP2C19 plays a minor role in the elimination of methadone (Dolophine).

Cardiovascular medications that are metabolized by CYP2C19 include the prodrug clopidogrel (Plavix), propranolol (Inderal), and cimetidine (Zantac). Proton-pump inhibitors such as omeprazole (Prilosec), esomeprazole (Nexium), lansoprazole (Prevacid), dexlansoprazole (Dexilant), and pantoprazole (Protonix) are major substrates of CYP2C19.

Inhibitors or inducers of CYP2C19 enzyme may modify its activity and change the patient's metabolizer status. This can result in drug-drug interactions when a drug substrate is prescribed with known CYP2C19 inhibitors or inducers.

Inhibitors

Some known CYP2C19 inhibitors include: fluconazole (Diflucan), fluvoxamine (Luvox), fluoxetine (Prozac), felbamate (Felbatol), ticlopidine (Ticlid), omeprazole (Prilosec), esomeprazole (Nexium), voriconazole (Vfend), armodafinil (Nuvigil), delavirdine (Rescriptor), modafinil (Provigil), oxcarbazepine (Trileptal), etravirine (Intelence), topiramate (Topamax), and moclobemide (Manerix).

Inducers

Some known CYP2C19 inducers include: artemether (Coartem), carbamazepine (Tegretol), efavirenz (Sustiva), enzalutamide (Xtandi), fosphenytoin (Cerebyx), primidone (Mysoline), phenobarbital, phenytoin (Dilantin), rifampin (Rifadin), and St. John's wort.

References

Preskorn SH. Clinically important differences in the pharmacokinetics of the ten newer "atypical" antipsychotics: Part 2. Metabolism and elimination. J Psychiatr Pract. 2012 Sep;18(5):361-8. 2: Preskorn SH. Clinically important differences in the pharmacokinetics of the ten newer "atypical" antipsychotics: part 1. J Psychiatr Pract. 2012 May;18(3):199-204. 3: Hicks et al. Clinical Pharmacogenetics Implementation Consortium Guideline for CYP2D6 and CYP2C19 Genotypes and Dosing of Tricyclic Antidepressants. Clin Pharmacol Ther. 2013 Jan 16;94(1):16-24. 4: Swen et al. Pharmacogenetics: from bench to byte--an update of guidelines. Clin Pharmacol Ther. 2011 May;89(5):662-73. 5: Wilffert et al. KNMP working group Pharmacogenetics. From evidence based medicine to mechanism based medicine. Reviewing the role of pharmacogenetics. Int J Clin Pharm. 2011 Feb;33(1):3-9. 6-Psychiatric Pharmacogenomics. David A. Mrazek. Publisher: Oxford University Press, USA; 1 edition (May 28, 2010)

CYP2D6

Clinical Utility

The cytochrome P450 2D6 (CYP2D6) is involved in the metabolism of 25% of clinically important medications. This enzyme is highly polymorphic: more than 100 different variants have been identified. The CYP2D6 assay identifies common variants that are associated with variability in CYP2D6 enzyme activity, which has important pharmacological and toxicological implications for antidepressants, antipsychotics, opioids, beta-blockers, and antiarrhythmics.

Assay Interpretation

CYP2D6 enzyme activity defines a normal or abnormal (intermediate, poor, or rapid) metabolizer status for a given individual. Commonly tested CYP2D6 variant alleles are classified into functional groups: Full or normal function (e.g., CYP2D6 *1, *2 and *35), reduced function (e.g., CYP2D6*9, *10, *148, *17, *29, and *41) and no function (e.g., CYP2D6 *3, *4, *5, *6, *7, *8, *11, *12, *14A, *15, *36 and *56). Increased CYP2D6 activity is found in individuals carrying multiple copies of functional alleles. CYP2D6 is subject to gene duplications, and these are denoted "XN", where N represents the number of CYP2D6 gene copies when available.

The genotype-phenotype relationship is established using a scoring system that assigns an activity value to every CYP2D6 allele, in order to assign a predicted phenotype. For a given genotype, the activity values of the constituent alleles are added together to calculate the CYP2D6 activity score. This activity score (AS) is then used to assign a predicted phenotype as follows: AS of 0 predicts a poor metabolizer, AS ranging between 1 and 2 predicts a normal (extensive) metabolizer, AS=0.5 predicts an intermediate metabolizer, and AS greater than 2 predicts a rapid (ultra-rapid) metabolizer. Fully functional alleles are assigned an activity value of 1, reduced function alleles have an activity value of 0.5, while non-functional alleles are assigned an activity value of 0.

The reference range for CYP2D6 metabolic status is a CYP2D6 *1/ *1 genotype, which is consistent with a normal metabolizer.

Clinical Implications

There is substantial evidence linking the CYP2D6 polymorphisms to variability in the pharmacological and safety profiles of the following psychotropics: desipramine (Norpramin), imipramine (Tofranil), amitriptyline (Elavil), nortriptyline (Pamelor), haloperidol (Haldol), trimipramine (Surmontil), venlafaxine (Effexor), doxepin (Silenor), aripiprazole (Abilify), atomoxetine (Strattera), duloxetine (Cymbalta), risperidone (Risperdal), clomipramine (Anafranil), and pimozide (Orap).

CYP2D6 polymorphisms have been shown to affect the pharmacological and safety profiles of the following analgesics: codeine, tramadol (Ultram), and hydrocodone (Vicodin). Codeine and tramadol are prodrugs that need to be activated by CYP2D6. Poor metabolizers are at high risk of therapy failure when given codeine or tramadol. On the other hand, rapid metabolizers may experience increased toxicity when given standard dosage of codeine or tramadol. Because CYP3A4 is also involved in the metabolism of oxycodone, patients with abnormal CYP2D6 activity may still experience adequate analgesia when taking this drug. CYP2D6 polymorphism has been shown to affect dihydrocodeine (Synalgos-DC) pharmacokinetics and can potentially alter the response to this drug.

Morphine, oxymorphone (Opana), hydromorphone (Dilaudid), butorphanol (Stadel), fentanyl (Duragesic), buprenorphine (Butrans), methadone (Dolophine), morphine (Avinza), and tapentadol (Nucynta) are not substrates of CYP2D6, and the patient's response to these drugs is not expected to be affected by polymorphisms in this enzyme.

Several important cardiovascular medications are metabolized by CYP2D6, and include: metoprolol (Lopressor), carvedilol (Coreg), flecainide (Tambocor), and propafenone (Rythmol).

Inhibitors of the CYP2D6 enzyme may modify its activity and change the patient's metabolizer status. This can result in drug-drug interactions when a drug substrate is prescribed with known CYP2D6 inhibitors. Although there are no known clinical inducers of CYP2D6, the pharmacokinetics of a drug substrate can be affected by inducers of other enzymes (such as CYP3A) that are involved in the metabolism of that drug.

Inhibitors

Some known strong and moderate CYP2D6 inhibitors include: abiraterone (Zytiga), bupropion (Wellbutrin), cobicistat (Stribild), fluoxetine (Prozac), quinidine (Quinidex), paroxetine (Paxil), cinacalcet (Sensipar), duloxetine (Cymbalta), terbinafine (Lamisil), tipranavir/ritonavir (Aptivus), mirabegron (Myrbetriq), peginterferon alfa-2b (Sylatron) and ecstasy.

Some known weak CYP2D6 inhibitors include: amiodarone (Cordarone), celecoxib (Celebrex), clobazam (Onfi), desvenlafaxine (Pristiq), diltiazem (Cardiazem), diphenhydramine (Benadryl), escitalopram (Lexapro), febuxostat (Uloric), gefitinib (Iressa), hydralazine (Apresoline), hydroxychloroquine (Plaquenil), imatinib (Gleevec), methadone (Dolophine), perphenazine (Trilafon), propafenone (Rythmol), ranitidine (Zantac), ritonavir (Norvir), sertraline (Zoloft), telithromycin (Ketek), verapamil (Isoptin, Cevera-HS), venlafaxine (Effexor), and Echinacea.

References

1- Swen et al. Pharmacogenetics: from bench to byte-an update of guidelines. Clin Pharmacol Ther. 2011 May;89(5):662-73. 2: Zhou SF. Polymorphism of human cytochrome P450 2D6 and its clinical significance: part II. Clin Pharmacokinet. 2009;48(12):761-804. 3: Zhou SF. Polymorphism of human cytochrome P450 2D6 and its clinical significance: Part I. Clin Pharmacokinet. 2009;48(11):689-723. 4: Preskorn SH. Clinically important differences in the pharmacokinetics of the ten newer "atypical" antipsychotics: Part 2. Metabolism and elimination. J Psychiatr Pract. 2012 Sep;18(5):361-8. 5: Preskorn SH. Clinically important differences in the pharmacokinetics of the ten newer "atypical" antipsychotics: part 1. J Psychiatr Pract. 2012 May;18(3):199-204. 6: D'Empaire et al. Antidepressant treatment and altered CYP2D6 activity: are pharmacokinetic variations clinically relevant? J Psychiatr Pract. 2011 Sep;17(5):330-9. 7: Hicks et al. Clinical Pharmacogenetics Implementation Consortium Guideline for CYP2D6 and CYP2C19 Genotypes and Dosing of Tricyclic Antidepressants. Clin Pharmacol Ther. 2013 Jan 16. 8: Gaedigk et al. The CYP2D6 activity score: translating genotype information into a qualitative measure of phenotype. Clin Pharmacol Ther. 2008 Feb;83(2):234-42. 9- Crews et al. Clinical Pharmacogenetics Implementation Consortium (CPIC) guidelines for codeine therapy in the context of cytochrome P450 2D6 (CYP2D6) genotype. Clin Pharmacol Ther. 2012 Feb;91(2):321-6. 10- Meyer et al. Absorption, distribution, metabolism and excretion pharmacogenomics of drugs of abuse. Pharmacogenomics. 2011 Feb;12(2):215-3

CYP3A4

Clinical Utility

The cytochrome P450 3A4 and 3A5 (CYP3A4 and CYP3A5) account for 40-80% of total CYP in human liver and intestine, respectively. Most importantly, CYP3A enzymes metabolize 50% of commonly used drugs. CYP3A4 and CYP3A5 enzymes have overlapping substrate specificity, and the contribution of CYP3A5 in the overall metabolism is smaller than the one for CYP3A4. The overall CYP3A metabolism status is expected to affect drugs that have a narrow therapeutic index.

Assay Interpretation

A limited number of variants identified within the CYP3A4 and CYP3A5 genes have been associated with significant alterations in enzyme activity and subsequent variability in therapeutic response. For CYP3A5, individuals with the less prevalent "normal metabolizer phenotype" may metabolize drugs faster than those with the more common "poor metabolizer phenotype". This may result in increased toxicity or loss of efficacy.

The CYP3A4*1 B variant is the most studied, and results in an enzyme with a moderately decreased activity. It occurs in 50% of African-Americans, 3-5% of Caucasians, and <1% of Asians. The CYP3A4*2, *3, *12, and *17 are also considered decreased activity alleles. Recently, the CYP3A4 *22 allele has been characterized as a decreased function allele that can be clinically relevant (associated with a decreased clearance of certain substrates). The genotype-phenotype relationship for CYP3A4 is not well established, and individuals are predicted to have a CYP3A4 normal or intermediate metabolic capacity.

The reference range for CYP3A4 metabolic status is CYP3A4 *1/ *1, which is consistent with a normal metabolizer.

The CYP3A5*3 variant results in an enzyme with no activity, and is the most common variant in the general population. The CYP3A5*3B, *6 and *7 are also null alleles resulting in no enzyme activity. The functional effects of the CYP3A5 alleles *2, *4, *5, *8, and *9 are not well established. The CYP3A5 *1 allele produces an active enzyme, and is found in 5% of Caucasians, 20% of Asians, and 15-50% of Africans. Individuals with two CYP3A5 inactive alleles are classified as poor metabolizers. Individuals carrying at least one copy of a CYP3A5 active allele are normal or intermediate metabolizers. CYP3A5 poor metabolizers represent 50% of Asians and 90% of Caucasians.

The reference range for CYP3A5 metabolic status is CYP3A5 *1/ *1, which is consistent with a normal metabolizer. This genotype is the least prevalent in Caucasians and Asians.

Clinical Implications

CYP3A4 and CYP3A5 genotypes can help identify patients with high or low overall CYP3A activity. Although these two enzymes metabolize many drugs, the response of only a few (such as narrow therapeutic index drugs) is expected to change significantly by genetic polymorphisms. Fentanyl (Duragesic) is a narrow therapeutic drug that is mainly metabolized by CYP3A. There is limited evidence suggesting that the response to this drug is altered in individuals with abnormal CYP3A activity.

The following drugs used in pain management and various psychiatric conditions are metabolized extensively by CYP3A: fentanyl (Duragesic), oxycodone (Oxycontin), buprenorphine (Suboxone), carbamazepine (Tegretol), quetiapine (Seroquel), ziprasidone (Geodon), alprazolam (Xanax), midazolam (Versed), triazolam (Halcion), nefazodone (Serzone), trazodone (Oleptro), vilazodone (Vibryd), zaleplon (Sonata), and zolpidem (Ambien). CYP3A contributes to a small extent in the elimination of methadone (Dolophine).

Within the major therapeutic classes used in cardiovascular conditions, the following drugs are substantially metabolized by CYP3A: atorvastatin (Lipitor), simvastatin (Zocor), lovastatin (Mevacor), nifedipine (Procardia), verapamil (Verelan), nicardipine (Cardena), felodipine (Plendil), nisoldipine (Sular), clopidogrel (Plavix), prasugrel (Effient), ticagrelor (Brilinta), cilostazol (Pletal), amiodarone (Cordarone), quinidine (Qualaquin), disopyramide (Norpace), losartan (Cozaar), rivaroxaban (Xarelto), and apixaban (Eliquis).

CYP3A metabolism is highly sensitive to inhibition and induction when a patient is taking multiple drugs. In this case, occurrence of drug-drug interactions can have profound effects on the pharmacokinetics, as well as the responses and safety profiles of many CYP3A drug substrates.

Genemarkers Gene Monographs

126 E. South Street | Kalamazoo | Michigan | 49007

Inhibitors

Some known strong CYP3A inhibitors include: ketoconazole (Nizoral), itraconazole (Sporanox), posaconazole (Noxatil), voriconazole (Vfend), clarithromycin (Biaxin), telithromycin (Ketek), troleandomycin (TAO), conivaptan (Vaprisol), nefazodone (Serzone), ritonavir (Norvir), saquinavir (Invirase), lopinavir, (Kaletra), nelfinavir (viracept), tipranavir (aptivus), boceprevir (Victrelis), telaprevir (Incivek), grapefruit juice (high dose) and idelalisib (Zydelig).

Some known moderate CYP3A inhibitors include: amprenavir (agenerase), aprepitant (Emend), atazanavir (Reyataz), darunavir (Prezista), fosamprenavir (Lexiva), erythromycin (Eryc), ciprofloxacin (Cipro), diltiazem (cardizem), verapamil (Isoptin, Covera-HS), fluconazole (Diflucan), imatinib (Gleevec), quinupristin/dalfopristin (Synercid), and grapefruit juice (low dose).

Some known weak CYP3A inhibitors include: amiodarone (Cordarone), amlodipine (Norvasc), atorvastatin (Lipitor), bicalutamide (Casodex), cilostazol (Pletal), fluvoxamine (Luvox), fluoxetine (Prozac), sertraline, cimetidine, ranitidine (Zantac), ranolazine (Ranexa), and ticagrelor (Brilinta).

Inducers

Some known strong CYP3A inducers include: carbamazepine (Tegretol), enzalutamide (Xtandi), fosphenytoin (Cerebyx), phenytoin (Dilantin), phenobarbital, primidone (Mysoline), rifampin (Rifadin), rifapentine (Priftin), and St. John's wort.

Some known moderate CYP3A inducers include: artemether (Coartem), bosentan (Tracleer), dabrafenib (Tatinlar), efavirenz (Sustiva), etravirine (Intence), modafinil (Provigil), nafcillin (Unipen), rifabutin (Mycobutin), and nevirapine (Viramune).

Some known weak CYP3A inducers include: fosamprenavir (Lexiva), aprepitant (Emend), clobazam (Onti), Echinacea, pioglitazone (Actos), dexamethasone (Decadron), oxcarbazepine (Trileptal), methylprednisolone (Medrol), and rutinamide (Banzel).

References

1- Metabolic Drug Interactions. RH Levy, KE Thummel, WF Trager. Publisher: Lippincott Williams & Wilkins (March 15, 2000). 2: Zhou et al. Polymorphism of human cytochrome P450 enzymes and its clinical impact. Drug Metab Rev. 2009;41(2):89-295 3- Isoherranen et al. The influence of CYP3A5 expression on the extent of hepatic CYP3A inhibition is substrate-dependent: an in vitro-in vivo evaluation. Drug Metab Dispos. 2008 Jan;36(1):146-54. 4- Williams et al. A significant drug-metabolizing role for CYP3A5? Drug Metab Dispos. 2003 Dec;31(12):1526-30. 5- Elens et al. A new functional CYP3A4 intron 6 polymorphism significantly affects tacrolimus pharmacokinetics in kidney transplant recipients. Clin Chem. 2011 Nov;57(11):1574-83. 6- Elens et al. Effect of a new functional CYP3A4 polymorphism on calcineurin inhibitors' dose requirements and trough blood levels in stable renal transplant patients. Pharmacogenomics. 2011 Oct;12(10):1383-96. 7- Lamba et al. Genetic contribution to variable human CYP3A-mediated metabolism. Adv Drug Deliv Rev. 2002 Nov 18;54(10):1271-94.

FACTOR II

Clinical Utility

Clotting Factor II, or prothrombin, is a vitamin K-dependent proenzyme that functions in the blood coagulation cascade. It is a precursor to thrombin, which converts fibrinogen into fibrin, which in turn strengthens a protective clot.

The prothrombin 20210G>A mutation in the Factor II gene results in increased levels of plasma prothrombin and a concurrent increased risk for thrombosis. Prothrombin-related thrombophilia is characterized by venous thromboembolism (VTE). This risk of thrombosis is also increased when mutations exist for other coagulation factors such as Factor V Leiden, or in presence of non-genetic risk factors such as obesity, injury, surgery, smoking, pregnancy, use of estrogen-containing contraceptives or replacement therapy. The clinical expression of Factor II thrombophilia is variable, and many individuals may never develop thrombosis, while others may experience venous thrombotic events or pregnancy complications.

Assay Interpretation

The Factor II thrombophilia is the second most common inherited risk factor for thrombosis. The Factor II 20210G>A mutation is associated with a hypercoagulable state. In the United States, the prevalence of this mutation is 1.1% in Caucasians and Hispanics and 0.3% in African-Americans. The prevalence of heterozygosity is 2%-5% in whites and 0%-0.3% in African-Americans. The prevalence of homozygosity is approximately one in 10,000.

The reference range for Factor II 20210G>A mutation is Factor II 20210GG.

Clinical Implications

The Factor II 20210G>A mutation is associated with an elevation of plasma prothrombin levels to about 30% above normal in heterozygotes and to 70% above normal in homozygotes. Heterozygotes are at a 2- to 5-fold increased risk of an initial VTE. The risk for VTE in Factor II 20210G>A homozygotes is not well defined, but is presumed to be higher than in 20210G>A heterozygotes. Factor II 20210G>A homozygotes tend to develop thrombosis more frequently and at a younger age. Individuals who are doubly heterozygotes for Factor V Leiden and Factor II 20210G>A have an estimated 20-fold increased risk when compared to individuals without either mutation, suggesting a multiplicative elevation in risk. Certain circumstantial factors can increase the risk of thrombosis, and include: travel, central venous catheter use, pregnancy, oral contraceptive use, hormone replacement therapy (HRT), selective estrogen receptor modulators (SERMs), organ transplantation, injury, age, and surgery.

Management Guidelines

Decisions regarding prophylactic anticoagulation should be based on a risk/benefit assessment. The first acute thrombosis should be treated according to standard guidelines. Following a first acute thrombotic event, long-term anticoagulation can be considered in homozygotes or in individuals with multiple thrombophilic disorders. In the absence of a history of thrombosis, anticoagulation is not routinely recommended for asymptomatic individuals who are heterozygotes. In asymptomatic individuals with a history of thrombosis and with one or two Factor II 20210G>A mutation(s), a short course of prophylactic anticoagulation may be considered in high-risk settings such as surgery, pregnancy, or prolonged immobilization. 2- Estrogen-containing contraceptives and replacement therapies: Heterozygote or homozygous women for Factor II 20210G>A mutation and with a history of VTE should avoid estrogen contraception and replacement therapy. Asymptomatic women who are heterozygote for Factor II 20210G>A mutation should be counseled on the risks of estrogen contraception and replacement therapy. Asymptomatic heterozygote women electing to use oral contraceptives should avoid third-generation formulations because of their higher thrombotic risk. Women homozygous for the Factor II 20210G>A mutation with or without prior VTE should avoid estrogen contraception and replacement therapy.

References

- 1- Gene Review: Prothrombin-Related Thrombophilia. Kujovich (2011) Available at <http://www.ncbi.nlm.nih.gov/books/NBK1148/> accessed on Mar 2013.
- 2- American College of Medical Genetics Consensus Statement on Factor V Leiden Mutation Testing. Grody et al. available at: (http://www.acmg.net/StaticContent/StaticPages/Factor_V.pdf) accessed on Mar 2013.
- 3- Evaluation of Genomic Applications in Practice and Prevention (EGAPP) Working Group. Recommendations from the EGAPP Working Group: routine testing for Factor V Leiden (R506Q) and prothrombin (20210G>A) mutations in adults with a history of idiopathic venous thromboembolism and their adult family members. Genet Med. 2011 Jan;13(1):67-76.
- 4- Segal et al. Predictive value of Factor V Leiden and Prothrombin G20210A in adults with venous thromboembolism and in family members of those with a mutation: a systematic review. JAMA. 2009 Jun 17;301(23):2472-85.

FACTOR V LEIDEN

Clinical Utility

The Factor V gene encodes the coagulation Factor V. In normal conditions, Factor V is inactivated during the clotting process by the activated protein C (APC). In subjects with Factor V Leiden thrombophilia, a mutation in the gene produces a Factor V that cannot be inactivated normally by APC. As a result, the clotting process remains active longer than usual, leading to more thrombin generation. This hypercoagulable state is also increased when other mutations exist in other coagulation factors such as Factor II, or in the presence of non-genetic risk factors such as obesity, injury, surgery, smoking, pregnancy, or use of estrogen-containing contraceptive or estrogen containing replacement therapy. The clinical expression of Factor V Leiden thrombophilia is variable. Many individuals may never develop thrombosis, while others may experience venous thrombotic events or pregnancy complications. Certain circumstantial factors can increase the risk of thrombosis, and include: travel, central venous catheter use, pregnancy, oral contraceptive use, hormone replacement therapy (HRT), selective estrogen receptor modulators (SERMs), organ transplantation, injury, age, and surgery. These factors are associated with the first thrombotic episode in at least 50% of individuals with a Factor V Leiden mutation.

Assay Interpretation

Factor V Leiden is the most common known inherited risk factor for thrombosis. Factor V Leiden refers to a base change (from G to A at position 1691) in the gene. As a result, Factor V is inactivated to a lesser extent and persists for longer in the circulation, leading to hypercoagulability. In the US, the frequency of the Factor V Leiden mutation varies by ethnicity, with about 5% of Caucasians, 2% of Hispanics, and 1% of African-Americans having one mutation. Only 1 in 5000 individuals have two Factor V Leiden mutations.

The reference range for Factor V Leiden mutation is Factor V 1691 GG.

Clinical Implications

About 1 in 1000 people in the U.S. experience a first venous thromboembolism (VTE) each year. VTE is caused by inherited and environmental factors, and while the Factor V Leiden mutation is present in only 15-20% of individuals with a first VTE, it is found in 50% of individuals with recurrent VTE or estrogen-related thrombosis. The risk for VTE is increased 3- to 8-fold in Factor V Leiden heterozygotes and 9- to 80-fold in homozygotes. This risk is increased further if other genetic or circumstantial factors are present. A heterozygote individual for both the Factor V Leiden and the Factor II (compound heterozygote) has an even greater risk of VTE (20-fold) than an individual with a mutation in only one factor. This illustrates the multiplicative effect of these two factors on overall thrombotic risk.

Management Guidelines

Decisions regarding prophylactic anticoagulation should be based on a risk/benefit assessment. The first acute thrombosis should be treated according to standard guidelines. Following a first acute thrombotic event, long-term anticoagulation can be considered in individuals homozygous for the Factor V Leiden mutation or with multiple thrombophilic disorders. In the absence of a history of thrombosis, anticoagulation is not routinely recommended for asymptomatic individuals who are heterozygote for the Factor V Leiden allele, because the 1- 3%/year risk for major bleeding from anticoagulation is greater than the estimated < 1%/year risk for thrombosis. In heterozygote individuals, a short course of prophylactic anticoagulation during exposure to hemostatic stresses may be considered. These situations include: surgery, pregnancy, and prolonged immobilization. 2- Estrogen-containing contraceptives and replacement therapies: women with one or two Factor V Leiden mutations and a history of VTE should avoid estrogen contraception and replacement therapy. Asymptomatic women who are heterozygote for Factor V Leiden should be counseled on the risks of estrogen contraception and replacement therapy. Asymptomatic women electing to use oral contraceptives should avoid third-generation formulations because of their higher thrombotic risk. Homozygote women with or without prior VTE should avoid estrogen contraception and replacement therapy.

References

- 1- Gene Review: factor V Leiden Thrombophilia. Kujovich (2010) Available at <http://www.ncbi.nlm.nih.gov/books/NBK1368/> accessed on Mar 2013.
- 2- American College of Medical Genetics Consensus Statement on Factor V Leiden Mutation Testing. Grady et al. available at: (http://www.acmg.net/StaticContent/StaticPages/Factor_V.pdf accessed on Mar 2013.
- 3- Rosendaal et al. Genetics of venous thrombosis. J Thromb Haemost. 2009 Jul;9 Suppl 1:301-4.
- 4- Bezemer et al. The value of family history as a risk indicator for venous thrombosis. Arch Intern Med. 2009 Mar 23;169(6):610-5.
- 5- Segal et al. Predictive value of Factor V Leiden and Prothrombin G20210A in adults with venous thromboembolism and in family members of those with a mutation: a systematic review. JAMA. 2009 Jun 17;301(23):2472-85.
- 6- Evaluation of Genomic Applications in Practice and Prevention (EGAPP) Working Group. Recommendations from the EGAPP Working Group: routine testing for Factor V Leiden (R506Q) and prothrombin (20210G>A) mutations in adults with a history of idiopathic venous thromboembolism and their adult family members. Genet Med. 2011 Jan;13(1):67-76

HTR2A

Clinical Utility

5-hydroxytryptamine (serotonin) receptor 2A (HTR2A) is a G protein-coupled receptor that encodes one of the receptors for serotonin. Serotonin is a neurotransmitter with many roles, including regulating mood, anxiety, feeding, and other behaviors, as well as dopamine and norepinephrine release in certain areas of the brain. HTR2A receptors are extensively distributed throughout the brain with high localization in the frontal and temporal regions. Preliminary studies indicate that some genetic variations in this gene are associated with susceptibility to schizophrenia and obsessive-compulsive disorder. They may also modulate the response to the antidepressant citalopram in patients with major depressive disorder (MDD).

Assay Interpretation

HTR2A variants have been studied with a primary objective of predicting response of the patients to anti-depressant therapy. Some of the most commonly tested variants of HTR2A are rs6311, rs6313 and rs7997012. Variants rs6311, also referred to as -1438G/A, is a guanine to adenine change and is located in the HTR2A promoter region. HTR2A with the adenine allele (A allele) is shown to have a greater expression when compared to the guanine allele (G allele). The other variant, rs6313 (102T/C), is a thymine to cytosine change. In samples of European ancestry, rs6311 is found to be in strong to complete linkage disequilibrium with the variant rs6313.

HTR2A rs7997012 is the other variant that is commonly tested. This variant is associated with a guanine to adenine change in intron 2 and is not in linkage disequilibrium with rs6311 or rs6313.

Clinical Implications

HTR2A genotyping assay may be used to determine the response to antidepressants. HTR2A rs6311 with the adenine allele (A allele) is shown to have greater expression when compared to the guanine allele (G allele). It has been reported that in patients with European ancestry, presence of the A allele of rs6311 (-1438G/A) and T allele of rs6313 (102C/T) is associated with a favorable response to clozapine. However, this association may not be consistent across patients with non-European ancestry.

HTR2A variant rs7997012 is associated with a differential response to citalopram. Patients who are homozygous for the A allele of rs7997012 respond better to citalopram when compared to the patients who are homozygous for the G allele.

These HTR2A variants have not yet been unequivocally validated as predictors of drug response. Additional studies are needed to establish their clinical impact.

References

- 1: McMahon FJ, Buervenich S, Charney D, Lipsky R, Rush AJ, Wilson AF, Sorant AJ, Papanicolaou GJ, Laje G, Fava M, Trivedi MH, Wisniewski SR, Manji H. Variation in the gene encoding the serotonin 2A receptor is associated with outcome of antidepressant treatment. *Am J Hum Genet.* 2006 May;78(5):804-14.
- 2: Parsons MJ, D'Souza UM, Arranz MJ, Kerwin RW, Makoff AJ. The -1438A/G polymorphism in the 5-hydroxytryptamine type 2A receptor gene affects promoter activity. *Biol Psychiatry.* 2004 Sep 15;56(6):406-10.
- 3: Arranz MJ, Munro J, Sham P, Kirov G, Murray RM, Collier DA, Kerwin RW. Meta-analysis of studies on genetic variation in 5-HT2A receptors and clozapine response. *Schizophr Res.* 1998 Jul 27;32(2):93-9. PubMed PMID: 9713904.
- 4: Arranz MJ, Kerwin RW. Neurotransmitter-related genes and antipsychotic response: pharmacogenetics meets psychiatric treatment. *Ann Med.* 2000 Mar;32(2):128-33.
- 5: Arranz MJ, Kerwin RW. Advances in the pharmacogenetic prediction of antipsychotic response. *Toxicology.* 2003 Oct 1;192(1):33-5. PubMed PMID: 14511901.
- 6: Arranz MJ, Collier DA, Munro J, Sham P, Kirov G, Sodhi M, Roberts G, Price J, Kerwin RW. Analysis of a structural polymorphism in the 5-HT2A receptor and clinical response to clozapine. *Neurosci Lett.* 1996 Oct 18;217(2-3):177-8.
- 7: Staddon S, Arranz MJ, Mancama D, Mata I, Kerwin RW. Clinical applications of pharmacogenetics in psychiatry. *Psychopharmacology (Berl).* 2002 Jun;162(1):18-23. Epub 2002 Apr 25.
- 8: Ellingrod VL, Lund BC, Miller D, Fleming F, Perry P, Holman TL, Bever-Stille K. 5-HT2A receptor promoter polymorphism, -1438G/A and negative symptom response to olanzapine in schizophrenia. *Psychopharmacol Bull.* 2003 Spring;37(2):109-12.
- 9: Lane HY, Lee CC, Liu YC, Chang WH. Pharmacogenetic studies of response to risperidone and other newer atypical antipsychotics. *Pharmacogenomics.* 2005 Mar;6(2):139-49.

HTR2C

Clinical Utility

HTR2C gene encodes a receptor that responds to the endogenous neurotransmitter serotonin. This serotonin receptor is a G protein-coupled receptor with wide distribution across the central and peripheral nervous systems in humans. It mediates excitatory neurotransmission via serotonin. Serotonin signaling regulates mood, anxiety, feeding, and many other behaviors, as well as dopamine and norepinephrine release in certain areas of the brain. Genetic variation in the HTR2C gene is known to be partially involved in pathogenesis of some psychiatric disorders and adverse effects of antipsychotic medications. The HTR2C assay identifies mutations that are associated with altered serotonin receptor expression and function, which has important pharmacological and toxicological implications for the use of antipsychotic medications, such as olanzapine, clozapine and risperidone.

Assay Interpretation

Many human polymorphisms in the HTR2C gene have been identified, of which a single nucleotide polymorphism (SNP rs3813929) is most significantly associated with a protective effect against the side effect of weight gain in patients taking olanzapine. HTR2C gene is located on the X chromosome in humans and SNP rs3813929 represents a C > T mutation in the regulatory region of this gene. As males have one copy of the HTR2C gene and in females one of the two copies of the gene is repressed, polymorphisms at this receptor can affect the two sexes to differing extent. Because of limited evidence, males with one copy of the C allele and females with one or two copies of the C allele are provisionally considered to have normal regulation of HTR2C expression and function. Males with one copy of the T allele and females with two copies of the T allele are provisionally considered to have altered regulation of HTR2C expression and function. The frequency of the minor T allele occurs in about 1% of European Caucasian and Asian populations, but is very rare in African populations.

Another single nucleotide polymorphism (SNP rs1414334) in the HTR2C gene is most significantly associated with a protective effect against the side effect of metabolic syndrome in patients taking clozapine or risperidone. SNP rs1414334 represents a C > G mutation in an intron region. Because of limited evidence, males with one copy of the C allele and females with one or two copies of the C allele are provisionally considered to have normal regulation of HTR2C expression and function. Males with one copy of the G allele and females with two copies of the G allele are provisionally considered to have altered regulation of HTR2C expression and function. The frequency of the G allele varies in different ethnic populations, and is often higher than the reference C allele. G allele occurs in about 45-65% of African populations, 80-90% of European Caucasian, and more than 90% of Asian populations.

Clinical Implications

The HTR2C assay identifies mutations that lead to functional variability in serotonin receptor, which is associated with clinical response to antipsychotics. In particular, genotype of SNP rs3813929 is significantly associated with the side effect of weight gain in patients who are taking olanzapine for psychiatric disorders including schizophrenia. Patients who have the rs3813929 TT genotype (female) or T genotype (male) may have a favorable response to olanzapine and a decreased risk of weight gain when compared to patients who have the rs3813929 CC genotype (female) or C genotype (male). Other genetic and clinical factors may also influence a patient's response to olanzapine.

Genotype of SNP rs1414334 is significantly associated with the side effect of metabolic syndrome in patients taking clozapine or risperidone. Patients who have the rs1414334 GG genotype (female) or G genotype (male) may have a favorable response to clozapine or risperidone, and a decreased risk of developing metabolic syndrome when compared to patients who have the rs1414334 CC genotype (female) or C genotype (male). Other genetic and clinical factors may also influence a patient's response to clozapine and risperidone.

In addition, HTR2C SNPs rs3813929 and rs1414334 may have other clinical implications in the treatment of psychiatric disorders such as depression, obsessive-compulsive disorder and anxiety-related conditions.

Both these HTR2C variants have not yet been unequivocally validated as predictors of drug response. Additional studies are needed to establish their clinical impact.

References

- Gregoor, J.G., et al., Polymorphisms of the LEP, LEPR and HTR2C gene: obesity and BMI change in patients using antipsychotic medication in a naturalistic setting. *Pharmacogenomics*, 2011. 12(6): p.919-23.
- Laika, B., et al., Pharmacogenetics and olanzapine treatment: CYP1A2*1 F and serotonergic polymorphisms influence therapeutic outcome. *Pharmacogenomics J*, 2010. 10(1): p. 20-9.
- Godlewska, B.R., et al., Olanzapine-induced weight gain is associated with the -759C/T and -697G/C polymorphisms of the HTR2C gene. *Pharmacogenomics J*, 2009. 9(4): p. 234-41.
- Ellingrod, V.L., et al., Weight gain associated with the -759C/T polymorphism of the 5HT2C receptor and olanzapine. *Am J Med Genet B Neuropsychiatr Genet*, 2005. 134B(1): p. 76-8.
- Risselada, A.J., et al., Association between HTR2C gene polymorphisms and the metabolic syndrome in patients using antipsychotics: a replication study. *Pharmacogenomics J*, 2012. 12(1): p. 62-7.
- Mulder, H., et al., HTR2C gene polymorphisms and the metabolic syndrome in patients with schizophrenia: a replication study. *J Clin Psychopharmacol*, 2009. 29(1): p. 16-20.
- Mulder, H., et al., The association between HTR2C gene polymorphisms and the metabolic syndrome in patients with schizophrenia. *J Clin Psychopharmacol*, 2007. 27(4): p. 338-43.

Genemarkers Gene Monographs

126 E. South Street | Kalamazoo | Michigan | 49007

MTHFR

Clinical Utility

Methylenetetrahydrofolate reductase (MTHFR) is involved in folate metabolism and is essential for the remethylation of homocysteine. Two common mutations in the MTHFR gene: 677C>T and 1298A> result in an enzyme with decreased activity, which is linked to increased plasma homocysteine levels (i.e., hyperhomocysteinemia). Mild to moderate hyperhomocysteinemia has been identified as a risk factor for venous thromboembolism and other cardiovascular diseases such as coronary heart disease and stroke. Other conditions in which hyperhomocysteinemia is found include recurrent pregnancy loss, placental infarction, and birth defects. However, the causal role of MTHFR mutations in these conditions is not well established.

Assay Interpretation

The approximate minor allele frequencies for most populations are 30-50% for the MTHFR 1298 A variant and 18-30% for the MTHFR 677 T variant. Heterozygotes and homozygotes for the MTHFR 677C>T mutation have 60% and 30% of normal MTHFR activity, respectively. Heterozygotes and homozygotes for the MTHFR 1298A>C mutation have 80% and 60% of normal MTHFR activity, respectively.

The reference ranges for both mutations of MTHFR are 677CC and 1298AA. This is consistent with a normal MTHFR activity.

Clinical Implications

The MTHFR assay provides information about potential causes of elevated homocysteine, and approaches for addressing it.

- Homozygosity for the MTHFR 677C>T mutation (individual with MTHFR 677 TT genotype) predisposes for hyperhomocysteinemia (especially during times of folate insufficiency) and an increase in premature cardiovascular disease. Measurement of total plasma homocysteine is informative in this case.
- Compound heterozygosity (individual with MTHFR 677 CT and MTHFR 1298AC genotypes) is not associated with an increase in plasma homocysteine level. Measurement of total plasma homocysteine is informative in this case.
- Homozygosity or heterozygosity for the MTHFR 1298A>C mutation alone (individual with MTHFR 1298AC or MTHFR 1298CC genotypes) does not increase homocysteine levels. Similarly, heterozygosity for the MTHFR 677C>T mutation alone (individuals with MTHFR 677 CT genotype) does not increase homocysteine levels.

Hyperhomocysteinemia related to MTHFR genetic mutations has been associated with neural tube defects, stillbirths, and recurrent pregnancy loss. However, because hyperhomocysteinemia is multifactorial, involving a combination of other genetic, physiologic, and environmental factors, the presence of MTHFR mutations in an individual should not be used alone to predict the risk of these conditions.

The response to methotrexate, a drug used in cancer and autoimmune diseases, is affected by the presence of MTHFR genetic mutations. Methotrexate intolerance is observed in individuals that are heterozygous or homozygous for the MTHFR 677C>T mutation.

References

1: van der Put. A second common mutation in the methylenetetrahydrofolate reductase gene: an additional risk factor for neural-tube defects? *Am J Hum Genet.* 1998 May;62(5):1044-51. 2: Lewis et al. Meta-analysis of MTHFR 677C>T polymorphism and coronary heart disease: does totality of evidence support causal role for homocysteine and preventive potential of folate? *BMJ.* 2005 Nov;331(7524): 1053. 3: Kluijtmans et al. Molecular genetic analysis in mild hyperhomocysteinemia: a common mutation in the methylenetetrahydrofolate reductase gene is a genetic risk factor for cardiovascular disease. *Am J Hum Genet.* 1996 Jan;58(1):35-41. 4: Hickey et al. ACMG Practice Guideline: lack of evidence for MTHFR polymorphism testing. *Genet Med.* 2013 Feb;15(2):153-6. 5: Grady et al. ACMG Factor V. Leiden Working Group. American College of Medical Genetics consensus statement on factor V Leiden mutation testing. *Genet Med.* 2001 Mar-Apr;3(2):139-48. 6: Gatt et al. Hyperhomocysteinemia and venous thrombosis. *Semin Hematol.* 2007 Apr;44(2):70-6. 7: De Mattia E, Toffoli G. C677T and A1298C MTHFR polymorphisms, a challenge for antifolate and fluoropyrimidine-based therapy personalisation. *Eur J Cancer.* 2009 May;45(8):1333-51. 8: Toffoli et al. Pharmacogenetic relevance of MTHFR polymorphisms. *Pharmacogenomics.* 2008 Sep;9(9):1195-206. 9: Clarke et al. MTHFR Studies Collaborative Group. Homocysteine and coronary heart disease: meta-analysis of MTHFR case-control studies, avoiding publication bias. *PLoS Med.* 2012 Feb;9(2):10. 10: Weisberg I, Tran P, Christensen B, Sibani S, Rozen R. A second genetic polymorphism in methylenetetrahydrofolate reductase (MTHFR) associated with decreased enzyme activity. *Mal Genet Metab.* 1998 Jul;64(3):169-72. 11: Weisberg et al. The 1298A>C polymorphism in methylenetetrahydrofolate reductase (MTHFR): in vitro expression and association with homocysteine. *Atherosclerosis.* 2001 Jun;156(2):409-15.

OPRM1

Clinical Utility

"Mu" opioid receptors are the most important site of action of opioid drugs. Single polymorphisms in the human mu-opioid receptor (OPRM1) have been investigated for their role in human nociception, opiate efficacy, and addiction.

Assay Interpretation

The variant mostly studied is a single substitution at position 118, from an adenine to a guanine (A118G). This variant reduces the OPRM1 receptor signaling efficiency induced by exogenous opioids. Reduced OPRM1 mRNA expression levels were observed in carriers of the G variant. The variant allele (G) is present in 7-15% of Caucasians, 1.5% of African-Americans, and up to 48.5% of Asians. The major interest of this particular SNP is due to its pharmacological and physiological consequences; however the exact mechanism by which the altered receptor influences opioid analgesia is still unresolved. The presence of the G allele seems to reduce the effect of exogenous agonists but increase the effects of exogenous antagonists.

The reference range for the A118G SNP is A118G AA, and is associated with a normal OPRM1 receptor signaling efficiency.

Clinical Implications

The presence of the G allele (A118G) seems to be associated with pain sensitivity as well as opioid dosage requirements. But only weak evidence of these associations is available to date. It is suggested that patients carrying the G allele report higher intensity pain. In terms of drug response, patients with the G allele have a favorable response to the anti-addictive drug naltrexone. Several studies conducted in post- surgical settings or in cancer analgesia showed that G allele carriers require slightly higher doses of morphine or fentanyl. This association still needs to be confirmed in larger studies and does not hold in other situations such as labor pain.

References

- 1: Wu et al. Polymorphism of the micro-opioid receptor gene (OPRM1 118A>G) affects fentanyl-induced analgesia during anesthesia and recovery. *Mal Diagn Ther.* 2009;13(5) :331-7.
- 2: Menon et al. The human μ -opioid receptor gene polymorphism (A118G) is associated with head pain severity in a clinical cohort of female migraine with aura patients. *J Headache Pain.* 2012Oct;13(7):513-9.
- 3: Olsen et al. Pain intensity the first year after lumbar disc herniation is associated with the A118G polymorphism in the opioid receptor mu 1 gene: evidence of a sex and genotype interaction. *J Neurosci.* 2012 Jul 18;32(29):9831-4.
- 4: Reyes-Gibby et al. Exploring joint effects of genes and the clinical efficacy of morphine for cancer pain: OPRM1 and COMT gene. *Pain.* 2007 Jul;130(1-2) :25-30.
- 5: Uitsch et al. Cross-sectional analysis of the influence of currently known pharmacogenetic modulators on opioid therapy in outpatient pain centers. *Pharmacogenet Genomics.* 2009 Jun;19(6):429-36.
- 6: Walter C, Uitsch J. Meta-analysis of the relevance of the OPRM1118A>G genetic variant for pain treatment. *Pain.* 2009 Dec;146(3):270-5.
- 7: Zhang et al. Association of human micro-opioid receptor gene polymorphism A118G with fentanyl analgesia consumption in Chinese gynaecological patients. *Anaesthesia.* 2010Feb;65(2):130-5.
- 8: Zhang et al. Study of the OPRM1 A118G genetic polymorphism associated with postoperative nausea and vomiting induced by fentanyl intravenous analgesia. *Minerva Anesthesiol.* 2011 Jan;77(1):33-9.
- 9: Oertel et al. The mu-opioid receptor gene polymorphism 118A>G depletes alfentanil-induced analgesia and protects against respiratory depression in homozygous carriers. *Pharmacogenet Genomics.* 2006 Sep;16(9):625-36.
- 10: Zwisler et al. Lack of Association of OPRM1 and ABCB1 Single-Nucleotide Polymorphisms to Oxycodone Response in Postoperative Pain. *J Clin Pharmacol.* 2011 Mar 24.
- 11: Klepstad et al. Influence from genetic variability on opioid use for cancer pain: a European genetic association study of 2294 cancer pain patients. *Pain.* 2011 May;152(5):1139-45.
- 12: Kadiiev E, et al. Role of pharmacogenetics in variable response to drugs: focus on opioids. *Expert Opin Drug Metab Toxicol.* 2008 Jan;4(1):77-91.
- 13: Vuilleumier et al. Pharmacogenomic considerations in opioid analgesia. *Pharmacogenomics Pers Med.* 2012;5:73-87.
- 14: Walter et al. μ -opioid receptor gene variant OPRM1 118 A>G: a summary of its molecular and clinical consequences for pain. *Pharmacogenomics.* 2013 Nov;14(15):1915-25.
- 15: Thorsell A. The μ -opioid receptor and treatment response to naltrexone. *Alcohol Alcohol.* 2013 Jul-Aug;48(4) :402-8.
- 16: Setiawan et al. Influence of the OPRM1 A118G polymorphism on alcohol-induced euphoria, risk for alcoholism and the clinical efficacy of naltrexone. *Pharmacogenomics.* 2012 Jul;13(10):1161-72.
- 17: Kranzler et al. Variation in OPRM1 moderates the effect of desire to drink on subsequent drinking and its attenuation by naltrexone treatment. *Addict Biol.* 2013 Jan;18(1):193-201.
- 18: Chamorro et al. Association of μ -opioid receptor (OPRM1) gene polymorphism with response to naltrexone in alcohol dependence : a systematic review and meta-analysis. *Addict Biol.* 2012 May;17(3):505-12.

Genemarkers Gene Monographs

126 E. South Street | Kalamazoo | Michigan | 49007

SLC0181

Clinical Utility

The SLC0181 gene encodes a liver-specific transporter involved in the removal of endogenous compounds (bile acids, bilirubin) and drugs such as statins from the blood to the liver. Some variants of the SLC0181 gene result in a low-functioning protein, which impairs statin clearance, and may lead to an increased risk of muscle pain, tenderness, or weakness, called myopathy. Certain medications can potentially inhibit SLC0181, causing clinically significant drug interactions.

Assay Interpretation

There are several variants of the SLC0181 that define over 15 alleles. One relatively common variant 521T>C (rs4149056) results in a decreased SLC0181 function, which affects the transport of drug substrates such as statins. This variant is present alone on the *5 allele and in presence with another variant (388A>G; rs2306283) on the *15 allele. Both alleles are low-activity alleles (reduced hepatic uptake), and have a combined frequency of 15-20% in Caucasians, 10-15% in Asians, and 2% in sub-Saharan Africans and African-Americans.

The reference range for the 521T>C mutation of SLC0181 is 521 TT. This is consistent with a normal SLC0181 transport function.

Clinical Implications

All statins are substrates of SLC0181, but the effects of SLC0181 genetic polymorphism differ between individual statins. The effect is the largest on simvastatin, and individuals with the 521T>C variant have increased levels of the active simvastatin form. The variant is strongly associated with simvastatin-induced myopathy (with or without CK elevation), especially with high-dose simvastatin therapy. More than 60% of the myopathy cases could be attributed to its presence. The clinical spectrum of statin-induced myopathy ranges from a mild and common myalgia to a life-threatening and rare rhabdomyolysis. Other known risk factors for statin-induced myopathy include a high-statin dose, interacting drugs that raise statin levels, age, hypothyroidism, and certain inherited muscle disorders.

At therapeutic doses, the apparent sensitivity levels of the five statins to the presence of the 521T>C variant are simvastatin>pitavastatin>atorvastatin>pravastatin>rosuvastatin. Carriers of the 521 T>C variant should avoid high-dose simvastatin therapy. These patients can take other statins, such as atorvastatin, pitavastatin, rosuvastatin, or pravastatin, but at reduced doses. Fluvastatin is not affected by the 521 T>C variant and could therefore be considered a suitable alternative.

Other drugs that are substrates of SLC0181 transporter include enalapril, olmesartan, valsartan, atrasentan, repaglinide, nateglinide, methotrexate, and bosentan. However, there is insufficient evidence documenting the impact of the 521 T>C variant on the systemic exposure and safety profile of these drugs.

Inhibitors

Inhibitors of SLC0181 transporter may alter its activity and result in increased levels of drug substrates. These include gemfibrozil, cyclosporine, clarithromycin, protease inhibitors, simeprevir, teriflunomide, boceprevir, telaprevir, and eltrombopag.

References

- 1: FDA Zocor Prescribing Label: <http://www.accessdata.fda.gov> 2: 1: Wilke et al. Clinical Pharmacogenomics Implementation Consortium (CPIC). The clinical pharmacogenomics implementation consortium: CPIC guideline for SLC0181 and simvastatin-induced myopathy. Clin Pharmacol Ther. 2012 Jul;92(1):112-7. 2: Feng et al. Individualized risk for statin-induced myopathy: current knowledge, emerging challenges and potential solutions. Pharmacogenomics. 2012 Apr;13(5):579-94. 3: Elsbey et al. Understanding the critical disposition pathways of statins to assess drug-drug interaction risk during drug development: it's not just about OATP1B1. Clin Pharmacol Ther. 2012 Nov;92(5):584-98. 4: SEARCH Collaborative Group, Link E. SLC0181 variants and statin-induced myopathy-a genome wide study. N Engl J Med. 2008 Aug 21;359(8):789-99. 5: Nies et al. Genetics is a major determinant of expression of the human hepatic uptake transporter OATP1B1, but not of OATP1B3 and OATP2B1. Genome Med. 2013 Jan 11;5(1):1. 6: Niemi M. Transporter pharmacogenetics and statin toxicity. Clin Pharmacol Ther. 2010 Jan;87(1):130-3. 7: Niemi et al. Organic anion transporting polypeptide 1B1: a genetically polymorphic transporter of major importance for hepatic drug uptake. Pharmacol Rev. 2011 Mar;63(1):157-81. 8: Neuvonen et al. Drug interactions with lipid-lowering drugs: mechanisms and clinical relevance. Clin Pharmacol Ther. 2006 Dec;80(6):565-81.

VKORC1

Clinical Utility

The Vitamin K epoxide reductase complex, subunit 1 (VKORC1) is the target of anticoagulants. This enzyme is the rate-limiting step in the vitamin K cycle. Mutations in the VKORC1 gene results in variable expression levels of the VKORC1 enzyme and altered sensitivities towards anticoagulants. VKORC1 genotype defines three levels of clinical phenotype: high, moderate, and low sensitivity phenotypes towards warfarin (a widely used anticoagulant). Therefore, VKORC1 variant testing is usually used in conjunction with CYP2C9 variant testing to optimize warfarin dosing and minimize the risks of bleeding or thrombotic complications.

Assay Interpretation

The clinically relevant variants in the VKORC1 gene are in strong linkage disequilibrium, meaning that some allele combinations occur more frequently than others. These combinations are referred to as haplotypes. The eight variants analyzed by the VKORC1 assay are used to define three haplotypes that are associated with different warfarin sensitivities, as shown in the following table.

Clinical Implications

The -1639G>A is the common variant seen in the Caucasian populations, and is believed to be the causative agent for the low-dose warfarin requirement phenotype. The G>A mutation results in a decreased expression of VKORC1. The 358C>T (found in 21% of African-Americans) and 3730G>A variants are associated with high warfarin dose requirements.

When CYP2C9 and VKORC1 genotypes are combined with other demographic (age, weight, height), clinical (disease, co-medications), and environmental (smoking) factors, they account for 50% of warfarin dose variation between individuals.

The FDA changed the warfarin label to help clinicians offer genotype-guided warfarin therapy for their patients.

References

1- Food and Drug Administration: Coumadin® Label accessed on Jan 2013. 2- Gage et al. Use of pharmacogenetic and clinical factors to predict the therapeutic dose of warfarin. Clin Pharmacol Ther. 2008 Sep;84(3):326-31 3- Schelleman et al. New genetic variant that might improve warfarin dose prediction in African Americans. Br J Clin Pharmacol. 2010 Sep;70(3):393-9 4- Johnson et al. Clinical Pharmacogenetics Implementation Consortium. Clinical Pharmacogenetics Implementation Consortium Guidelines for CYP2C9 and VKORC1 genotypes and warfarin dosing. Clin Pharmacol Ther. 2011 Oct;90(4):625-9. 5- Klein et al. Estimation of the warfarin dose with clinical and pharmacogenetic data. International Warfarin Pharmacogenetics Consortium. N Engl J Med. 2009 Feb 19;360(8):753-64 6-Rieder et al. Effect of VKORC1 haplotypes on transcriptional regulation and warfarin dose. N Engl J Med. 2005 Jun 2;352(22):2285-93.